



Diversity Analysis of Dairy and Nondairy *Lactococcus lactis* Isolates, Using a Novel Multilocus Sequence Analysis Scheme and (GTG)₅-PCR Fingerprinting[†]

Jan L. W. Rademaker,¹ Hélène Herbet,¹ Marjo J. C. Starrenburg,¹ Sabri M. Naser,² Dirk Gevers,² William J. Kelly,³ Jeroen Hugenholtz,¹ Jean Swings,² and Johan E. T. van Hylckama Vlieg^{1*}

*NIZO food research, Kluyver Centre for Genomics of Industrial Fermentation, P.O. Box 20, 6710 BA Ede, The Netherlands*¹;
*Laboratory of Microbiology and BCCM/LMG Bacteria Collection, Ghent University, K. L. Ledeganckstraat 35, B-9000 Ghent, Belgium*²; and *AgResearch Ltd., Grasslands Research Centre, Private Bag 11008, Palmerston North, New Zealand*³

Received 7 May 2007/Accepted 10 September 2007

The diversity of a collection of 102 *lactococcus* isolates including 91 *Lactococcus lactis* isolates of dairy and nondairy origin was explored using partial small subunit rRNA gene sequence analysis and limited phenotypic analyses. A subset of 89 strains of *L. lactis* subsp. *cremoris* and *L. lactis* subsp. *lactis* isolates was further analyzed by (GTG)₅-PCR fingerprinting and a novel multilocus sequence analysis (MLSA) scheme. Two major genomic lineages within *L. lactis* were found. The *L. lactis* subsp. *cremoris* type-strain-like genotype lineage included both *L. lactis* subsp. *cremoris* and *L. lactis* subsp. *lactis* isolates. The other major lineage, with a *L. lactis* subsp. *lactis* type-strain-like genotype, comprised *L. lactis* subsp. *lactis* isolates only. A novel third genomic lineage represented two *L. lactis* subsp. *lactis* isolates of nondairy origin. The genomic lineages deviate from the subspecific classification of *L. lactis* that is based on a few phenotypic traits only. MLSA of six partial genes (*atpA*, encoding ATP synthase alpha subunit; *pheS*, encoding phenylalanine tRNA synthetase; *rpoA*, encoding RNA polymerase alpha chain; *bcaT*, encoding branched chain amino acid aminotransferase; *pepN*, encoding aminopeptidase N; and *pepX*, encoding X-prolyl dipeptidyl peptidase) revealed 363 polymorphic sites (total length, 1,970 bases) among 89 *L. lactis* subsp. *cremoris* and *L. lactis* subsp. *lactis* isolates with unique sequence types for most isolates. This allowed high-resolution cluster analysis in which dairy isolates form subclusters of limited diversity within the genomic lineages. The *pheS* DNA sequence analysis yielded two genetic groups dissimilar to the other genotyping analysis-based lineages, indicating a disparate acquisition route for this gene.

Lactococcus lactis is the primary constituent of many industrial and artisanal starter cultures used to ferment dairy products, especially hard and semihard cheeses. These starter cultures play a key role in determining shelf-life, preservation, and organoleptic quality and thus influence the quality and safety of these fermented products (35). As a result of the industrial importance of *L. lactis*, it has been the subject of numerous studies, which have resulted in detailed knowledge of its physiology and molecular biology. Moreover, due to the availability of a vast molecular toolbox and three whole genome sequences, of strains IL-1403, SK11, and MG1363 (3, 22, 40), *L. lactis* has gained a strong position as a model organism for low-GC gram-positive bacteria (20).

The model strains that are used in most studies almost exclusively originate from dairy fermentations. Isolates from (fermented) plant material have been only poorly characterized. However, in recent years there has been growing interest in plant isolates as several examples indicate phenotypes of industrial interest such as a unique flavor-forming potential or the production of bacteriocins with a broad mode of action (1, 17, 18).

Historically, three different industrial phenotypes have been

recognized, those of *L. lactis* subsp. *lactis*, *L. lactis* subsp. *cremoris*, and *L. lactis* subsp. *lactis* biovar diacetylactis (34). These two subspecies and one biovar are taxonomically differentiated by only a few phenotypic characteristics (Table 1). *L. lactis* subsp. *cremoris* is characterized by the inability to produce ammonia from arginine and by low tolerance to elevated temperatures and salt concentrations. *L. lactis* subsp. *lactis* produces ammonia from arginine and is tolerant to 40°C and 4% NaCl (34). Some *L. lactis* subsp. *lactis* strains are able to ferment citrate and to produce the flavor compound diacetyl, and these are referred to as *L. lactis* subsp. *lactis* biovar diacetylactis strains (34).

Numerous studies involving DNA-DNA hybridization, small subunit (SSU) rRNA gene sequences, and PCR fingerprint analyses show two genomic lineages (7, 23, 28, 31). Additionally, subspecific probes have been proposed and applied for the identification of environmental isolates (18, 32). This started an increasingly confusing period of subspecies classification and identification of *L. lactis* strains. For example, two different genotypes recognized by SSU rRNA sequence differences were referred to as *L. lactis* subsp. *lactis* and *L. lactis* subsp. *cremoris* (31), but several groups have recognized that phenotypic and genotypic groupings do not match (16, 27, 32, 38). An *L. lactis* subsp. *cremoris*-like genotype, i.e., a genotype similar to the genotype of the *L. lactis* subsp. *cremoris* type strain, may have the *L. lactis* subsp. *lactis*

* Corresponding author. Mailing address: NIZO food research, P.O. Box 20, 6710 BA Ede, The Netherlands. Phone: 31 318 659 511. Fax: 31 318 650 400. E-mail: Johan.van.Hylckama.Vlieg@nizo.nl.

† Published ahead of print on 21 September 2007.

TABLE 1. Phenotypes and genotypes of *Lactococcus* species and subspecies and biovar diacetylactis

Organism	Phenotype ^a							Genotype of cluster ^b		
	Growth at 40°C	Growth with 4% NaCl	Citrate utilization	Arginine dehydrolase	Maltose ^c	Lactose and galactose ^c	Ribose ^c	SSU rRNA gene and (GTG) ₅ fp	<i>L. lactis</i> subsp. <i>lactis</i> type-strain-like (n)	<i>L. lactis</i> subsp. <i>cremoris</i> type-strain-like (n)
<i>L. lactis</i> subsp. <i>lactis</i>	+(−)	+	−	+	+	+	+	1A, 1B, 1C	1A (60), 1C (2)	1B (9)
<i>L. lactis</i> subsp. <i>lactis</i> biovar diacetylactis	+(−)	+	+	+	+	+	+	1A	1A (7)	
<i>L. lactis</i> subsp. <i>cremoris</i>	−	−	−	−	−	+	−	1A, 1B		1B (11)
<i>L. lactis</i> subsp. <i>hordniae</i>	−	−	−	+	−	−	−	1A	NA	NA
<i>L. raffinolactis</i>	−	−	−	− (++)	+	+	− (++)	3	NA	NA
<i>L. plantarum</i>	V	+	−	−	+	−	−	5	NA	NA
<i>L. garvieae</i>	+	+	−	+	+(−)	+	+	2	NA	NA
<i>L. piscium</i>	−	NA	NA	−	+	+	−	4	NA	NA

^a Adapted from Schleifer et al. and Holt et al. (13, 34). V, variable; NA, not available/applicable; +, positive; −, negative; +(−), most positive but occasional negative; − (++)+, most negative but occasional positive.

^b Data are from the present study. n, number of isolates in this study; NA, not available/applicable; fp, fingerprinting.

^c Results indicate whether acid is produced from this substance.

phenotype, with strain MG1363 a well-known representative of this group (38). Other investigators postulated that lactococci can be divided into three to five major groups with an unusual taxonomic structure (16, 36). In this structure, two genetically distinct groups of strains show indistinguishable phenotypes while, conversely, two distinct phenotypic groups are genetically homologous.

More recently, microbes have been characterized by DNA sequence analysis of several housekeeping genes or other protein-coding gene sequences in an approach called multilocus sequence analysis (MLSA) or typing (21). MLSA methods have brought a new dimension to the elucidation of genomic relatedness at inter- and intraspecific levels, providing microbiologists with tools to search for phylogenetic markers independent of ribosomal DNA genes, and are most promising in obtaining biologically meaningful and functional groupings (21). Several studies have been published describing the application of MLSA to estimate the relative contribution of recombination and (point) mutation to clonal diversification in order to study phylogeny and microbial diversity in specific ecological niches (4, 14, 21). So far, MLSA and multilocus sequence typing methods have been used primarily in epidemiology including the population analysis of food-related pathogens like *Salmonella*, *Listeria monocytogenes*, or *Bacillus cereus* (12, 33, 37). Most recently, a method was reported for *Lactobacillus plantarum* (6).

In the present study genotypic and the traditionally used differentiating phenotypic characteristics (Table 1) are determined to assess the diversity of 102 lactococcus isolates including a subset of 89 strains of *L. lactis* subsp. *cremoris* and *L. lactis* subsp. *lactis* isolates from dairy and nondairy (plant) origin. Additionally, the development and application of a novel MLSA scheme targeting housekeeping and flavor-related genes for diversity analysis within *L. lactis* is reported. The scheme shows the existence of three separate lineages. Additionally, diversity was studied using SSU rRNA gene sequence analysis and repetitive element-based PCR genomic fingerprint analysis (30).

MATERIALS AND METHODS

Bacterial isolates and medium. A total of 103 isolates were subjected to genotypic analysis (Fig. 1). The collection included 102 lactococcus isolates representing *L. garvieae*, *L. piscium*, *L. plantarum*, *L. raffinolactis*, *L. lactis* subsp. *hordniae*, *L. lactis* subsp. *cremoris*, and *L. lactis* subsp. *lactis*. Isolates from the latter two subspecies were labeled dairy or nondairy according to their sources of isolation, and care was taken to maximally cover the diversity within the species. Strains were maintained in M17 broth (Oxoid Ltd., Basingstoke, Hampshire, England) with 0.5% glucose (wt/vol) as a carbon source (GM17). Serial transfer was minimized to prevent the occurrence of mutations as a result of adaptation to laboratory medium and conditions. Additionally, *Enterococcus casseliflavus* NIZO 2193 was used in this study as an outgroup in DNA sequence analysis.

Genotypic diversity. (i) **DNA extraction.** Bacterial DNA was extracted using an InstaGene Matrix (Bio-Rad, Hercules, CA) according to the manufacturer's instructions.

(ii) **Genomic fingerprinting and computer-assisted analysis.** (GTG)₅-PCR genomic fingerprints were obtained as described previously (11, 30). Computer-assisted analysis of the genomic fingerprints was performed using BioNumerics software, version 4.50 (Applied Maths, Saint-Martens-Latem, Belgium), as described previously (29). Briefly, the similarity between pairs of genomic fingerprints was calculated using the product-moment correlation coefficient (r value), applied to the whole densitometric curve of the gel tracks. Cluster analysis of the pairwise similarity values was performed using the unweighted-pair group method using arithmetic averages (UPGMA) algorithm.

(iii) **Selection of loci for sequence analysis and primer design.** Multiple protein-coding loci, both housekeeping genes and functional genes, were evaluated for their applicability in assessing the diversity and relationships of different isolates in an MLSA approach. Criteria used were the following: (a) broad distribution among whole genome sequences of lactic acid bacteria ([LAB] 14 completed genomes and 10 draft genome sequences were included), (b) presence in only one copy per organism, and (c) mutually unlinked in location. The final selection included three housekeeping genes: *atpA*, encoding ATP synthase alpha subunit; *pheS*, encoding phenylalanine tRNA synthetase; and *rpoA*, encoding RNA polymerase alpha chain. Three functional genes involved in production of flavor compounds during fermentation were also included: *bcaT*, encoding branched chain amino acid aminotransferase; *pepN*, encoding aminopeptidase N; and *pepX*, encoding X-prolyl dipeptidyl peptidase. Multiple sequence alignments of these genes from all LAB and phylogenetic analyses showed that conserved regions, suitable for primer design, were not always present among all included LAB sequences. This rendered the construction of universal LAB primers unfeasible. Therefore, primers were designed on the conserved regions within the lactococcal genomes and extended with related genomes where possible, as long as two suitable conserved regions remained. With respect to the housekeeping genes, sequences from the following genomes were included: *Enterococcus faecalis* (V583), *Lactobacillus plantarum* (WCFS1), *L. lactis* subsp. *lactis* (IL-1403), *L. lactis* subsp. *cremoris* (SK11), *Streptococcus pneumoniae* (TIGR4 and R6), *Streptococcus agalactiae* (NEM316 and 2603 V/R), *Streptococcus pyogenes*




TABLE 2. Primers and annealing temperatures used for gene-specific PCR amplification and sequencing of *L. lactis* isolates

Gene	Forward primer ^a	Reverse primer ^a	Annealing temp (°C)
<i>rpoA</i>	5'-ATGATYGARTTTGAAAAACC-3'	5'-ACHGTRTRATDCCDGRCG-3'	46
<i>atpA</i>	5'-TAYRTYGGKGAYGGDATYGC-3'	5'-CCRCGRTTHARYTTHGCYTG-3'	50
<i>bcaT</i>	5'-TTTKSHRTGCCDGTVGG-3'	5'-GGWCCHACTTCYGTYTC-3'	46
<i>pepN</i>	5'-ATKCTTAYGCWGAYRTYGT-3'	5'-TTKCTTCAAGSMAWGSCC-3'	50
<i>pepX</i>	5'-TTTGGGTTGAAAGTCCAGT-3'	5'-CCAAGAAGAAATTCCAGC-3'	46
<i>pheS</i>	5'-CAYCCNGCHCGYGYATGC-3'	5'-CCWARVCCRAARGCAAARCC-3'	50
SSU rRNA	5'-GCGCGTGCCTAATACATGC-3'	5'-ATCTACGCATTTCACCGCTAC-3'	50

^a Y, R, H, D, K, Y, W, M, and N are degenerate nucleotides according to the degenerate nucleotide alphabet: R for A or G; M for A or C; Y for C or T; K for G or T; W for A or T; N for A, G, C or T; S for G or C; H for A, C or T; and D for A or G or T.

(MGAS8232, SSI-1, MGAS315, and SF370), and *Streptococcus mutans* (UA159). For *bcaT*, sequences from *E. faecalis* (V583), *L. lactis* subsp. *lactis* (IL-1403), *L. lactis* subsp. *cremoris* (SK11), *S. pneumoniae* (TIGR4 and R6), *S. agalactiae* (NEM316 and 2603 VR), *S. pyogenes* (MGAS8232, SSI-1, MGAS315, and SF370), and *S. mutans* (UA159) were included; for *pepN* both lactococcus genomes, *Lactobacillus johnsonii* (NCC 533), and *Lactobacillus gasseri* were included, whereas for *pepX* only the lactococcus genomes were used. Primer design was performed by standard procedures using Kodon, version 2.04, software (Applied Maths).

(iv) **Multilocus and SSU rRNA gene sequence analysis.** DNA sequence analysis was performed of approximately 400-bp intragenic regions of the *atpA*, *pheS*, *rpoA*, *bcaT*, *pepN*, and *pepXP* genes and the SSU rRNA gene. The primer combinations (Proligo, Paris, France) for the primary PCR and cycle sequencing and the annealing temperatures used are listed in Table 2. The amplification of the gene fragments was carried out for 30 cycles in a 50- μ l mixture containing each deoxynucleoside triphosphate at a concentration of 2.5 mM, 250 ng of each primer, 1 U of *Taq* polymerase (Applied Biosystems, Nieuwerkerk aan de IJssel, The Netherlands), and 5.0 μ l of genomic DNA. After incubation for 3 min at 95°C and for 15 s at 94°C, samples were subjected to 30 cycles of 30 s at the annealing temperature (Table 2), followed by 1 min 45 s at 72°C; the reaction was completed by 4 min at 72°C and kept at 4°C using a GeneAmp PCR System 9600 thermocycler (Applied Biosystems). Presence, size, and approximate quantity of the resulting PCR product were analyzed using a 1% agarose gel with 0.5 μ g of ethidium bromide per ml and a low DNA mass ladder (E-Gel; Invitrogen). The obtained PCR products were purified using a GFX PCR kit (Amersham Biosciences, Roosendaal, The Netherlands) according to the manufacturer's instructions. The forward primer of the primary PCR was also applied in cycle sequencing using an ABI PRISM BigDye Terminator (version 3.1) cycle sequencing kit (Applied Biosystems) according to the manufacturer's protocol. Briefly, 2.5 μ l of the primary PCR and 3 pmol of the forward primer were used in a 20- μ l total sequencing reaction mixture. Cycle sequencing started with 1 min at 96°C followed by 25 cycles of 96°C for 10 s and 50°C for 5 s, with a final extension for 4 min at 60°C. Alternatively, 46°C for 5 s was used for annealing when no PCR product was obtained at 50°C, but still no specific amplicons were obtained for limited genes and strains, as reported below. Reaction products were purified using Autoseq G-50 columns (Amersham Biosciences) according to the manufacturer's instructions. After denaturation, 10 μ l of the purified template was mixed with 30 μ l of Hi-Di (Applied Biosystems) and incubated for 5 min at 95°C. Electrophoresis was performed using an ABI Prism 310 Genetic Analyser, and

the sequences were captured using GeneScan analysis, version 6.7 (Applied Biosystems).

(v) **Computer-assisted analysis of DNA sequence data.** Single forward sequences were trimmed, aligned, and analyzed using BioNumerics, version 4.50, software. For each gene fragment, sequences were compared using neighbor-joining cluster analysis based on global alignment similarities. *E. casseliflavus* NIZO 2193 was used as outgroup in SSU rRNA gene sequence cluster analysis. Bootstrap analysis was applied with 500 simulations with single-sequence cluster analyses. Additionally, a composite data set was defined averaging the similarity matrices of five gene sequence analyses.


Phenotypic diversity. (i) **Growth tests.** For screening purposes isolates were grown at 30°C in 96-well microplates using 250 μ l of GM17 medium. Overnight cultures of isolates were incubated in quadruplicate, and each plate contained strain ML3 (NIZO 643) as reference and no inoculum as a negative control. The following test media and conditions were evaluated: GM17 medium at 37 and 45°C; GM17 medium containing 4% NaCl at 30°C. When the test medium was not GM17, strains were subcultured twice to prevent false positives due to carryover from GM17 medium. Growth was analyzed by determining the turbidity at 600 nm and scored as negative (0), weak (1), or positive (2). Growth was scored after 24 h.

(ii) **Fermentation tests.** For the determination of arginine hydrolyase activity, isolates were grown in 96-well microplates in arginine broth as described by Niven et al. (26) but supplemented with 5 g/liter peptone, 250 mg/liter $MgSO_4 \cdot 7H_2O$, and 0.5 g/liter ascorbic acid. Growth was scored after 48 h by determining the turbidity at 600 nm. Cells were removed by centrifugation, and 10 μ l of the resulting supernatant was mixed with reagent according to Niven et al. (5 g of KL, 5 g of $HgCl_2$, 4 g of NaOH, and 100 ml of H_2O , filtered) (26), and arginine hydrolyzing activity was scored as negative (0), weak (1), or strong (2), judged by the intensity of the orange color. Strain LMG 8505 (NIZO 2419) did not grow on the medium used by Niven et al. (26), and consequently no arginine hydrolyase activity was determined.

The ability to ferment citrate was measured with whey permeate calcium citrate Casitone agar (WPCCA) as described by Galesloot et al. (8) but with milk replaced by whey permeate.

The fermentation of maltose, lactose, galactose, and ribose was determined with the API 50 CHL assay (BioMérieux, l'Etoile, France). Strains were cultivated overnight in GM17 medium at 30°C, after which the culture was washed and resuspended in M17 with 1/4 sodium β -glycerol phosphate buffer (4.75 g/liter). Fermentation activity was determined after 48 h of incubation at 30°C.

FIG. 1. Neighbor-joining cluster analysis of the partial SSU rRNA gene sequence (*Escherichia coli* positions 41 to 705) of 102 *Lactococcus* isolates with *E. casseliflavus* as the outgroup. Bootstrap percentages (≥ 50) after 500 simulations are shown. All *L. lactis* subsp. *lactis* isolates and *L. garvieae* showed growth at 37°C; other isolates did not (data not shown). Further phenotypes characteristic for *L. lactis* subsp. *cremoris* and *L. lactis* subsp. *lactis*, according to Schleifer et al. (37), are indicated and include the production of ammonia from arginine, acid from maltose, and salt tolerance at 4% NaCl. Additionally, citrate production on WPCCA medium is specified for differentiation of biovar diacetylactis (data not shown). Isolates obtained from dairy substrates are indicated (■). Five distinct species clusters (1 to 5) support the *Lactococcus* classification. The *L. lactis* isolate cluster 1 includes three subgroups; one major subgroup, 1B, comprises 11 *L. lactis* subsp. *cremoris* and 9 *L. lactis* subsp. *lactis* isolates, whereas the other major subgroup, 1A, includes 67 *L. lactis* subsp. *lactis* and 2 *L. lactis* subsp. *hordniae* isolates. Finally, two *L. lactis* subsp. *lactis* strains form a distinct minor subcluster (1C). Culture references: LMG, Laboratorium voor Microbiologie, University of Ghent, Belgium; ATCC: American Type Culture Collection, Manassas, VA; DSM(Z), Deutsche Sammlung von Mikroorganismen und Zellkulturen, Braunschweig, Germany; NCFB, NCDO National Collection of Food Bacteria; National Collections of Industrial and Marine Bacteria Ltd., Aberdeen, Aberdeen, Scotland, United Kingdom. KF and KW strains originate from Kelly's laboratory, Palmerston North, New Zealand; other strains are from the NIZO culture collection.



using the recommendations of the manufacturer. The activity was scored using a seven-step incremental scale from negative (0) to positive (6).

Nucleotide sequence accession numbers. The sequences generated in this study are available from GenBank/EMBL databases under the accession numbers EU091374 to EU091475 (SSU rRNA partial gene sequences), EU111131 to EU111178 (*rpoA* partial gene sequences), EU111220 to EU111267 (*pheS* partial gene sequences), EU111568 to EU111615 (*atpA* partial gene sequences), EU111309 to EU111354 (*pepX* partial gene sequences), EU111396 to EU111442 (*pepN* partial gene sequences), and EU111483 to EU111529 (*bcaT* partial gene sequences).

RESULTS

Taxonomic framework based on SSU rRNA gene sequence and phenotypic analysis. Cluster analysis of partial SSU rRNA gene sequences of all isolates revealed five distinct species clusters (Fig. 1, clusters 1 to 5) that support the *Lactococcus* species classification of *L. lactis*, *L. garvieae*, *L. raffinolactis*, *L. piscium*, and *L. plantarum*. Additionally, all strains were screened using the phenotypic tests as proposed by Schleifer et al. (34) for the subspecific classification of *L. lactis* subsp. *cremoris* and *L. lactis* subsp. *lactis*. These tests include the production of ammonia from arginine and acid from maltose as well as salt tolerance at 4% NaCl (Table 1), and the results are depicted in Fig. 1. Eleven *L. lactis* isolates showed the inability to grow at NaCl concentrations of more than 4% at 37°C and did not form acid from maltose and ammonia from arginine, the discriminating phenotypic traits for *L. lactis* subsp. *cremoris* (Fig. 1). Furthermore, citrate production is specified for differentiation of biovar diacetyl-lactis, as indicated by the WPCCA column in Fig. 1. The *L. lactis* cluster (Fig. 1, cluster 1) includes two major subgroups: subgroup B comprises 11 *L. lactis* subsp. *cremoris* isolates (unable to grow at 37°C) and in addition 9 *L. lactis* subsp. *lactis* isolates (able to grow at 37°C), whereas subgroup A includes the other 67 *L. lactis* subsp. *lactis* isolates (able to grow at 37°C) and 2 *L. lactis* subsp. *hordniae* isolates (unable to grow at 37°C). The *L. lactis* subsp. *hordniae* sequences differ by one nucleotide (underlined) in the V1 region, having a T in AGTGGGGGA instead of the C in AGCGG

FIG. 2. Comparative diversity analysis of 89 *L. lactis* isolates. Neighbor-joining cluster analysis of the partial SSU rRNA gene sequence (*Escherichia coli* positions 41 to 705), a five-locus MLSA based on a composite data set of DNA sequences of ~0.4 kb of the intragenic regions of housekeeping genes *atpA* and *rpoA* and flavor-related genes *bcaT*, *pepN*, and *pepXP* and the single partial *pheS* sequence. Bootstrap percentages (≥ 50) after 500 simulations are shown for the single sequence analyses. Strains with the *L. lactis* subsp. *lactis* type-strain-like genotype [SSU rRNA gene sequence and (GTG)₅-PCR fingerprint type] are represented with gray bars. Strains with an *L. lactis* subsp. *cremoris* type-strain-like genotype [SSU rRNA gene sequence and (GTG)₅ PCR fingerprint type] are represented with white bars. Isolates obtained from dairy substrates are indicated (■). The results of the partial SSU rRNA gene sequence and five-locus MLSA cluster analysis allow the discrimination of two major genomic lineages designated the *L. lactis* subsp. *cremoris* and *L. lactis* subsp. *lactis* type-strain-like genotypes, respectively. Strains P7304 and P7266 (boxed) grouped separately from the two major lineages. No sequences of *pepN* for isolates KF7 and KF 178, of *pepXP* for strains P7266 and P7304, and of *bcaT* for isolates P7266, KF 178, KW2, and AM3 are included in the composite five-locus MLSA cluster analysis presented since they were not obtained experimentally (see explanation in text).




FIG. 2—Continued.

TABLE 3. Genetic diversity at seven loci based on 76 *L. lactis* isolates included in Fig. 2 and 3 for which sequences are available for all loci^a

Locus	Gene length (bp)	Amplified fragment length (bp)	Analyzed fragment length (bp)	No. of polymorphic sites	% of variable nucleotide sites	No. of alleles	Avg G+C content (%)	dN/dS ratio ^b
<i>atpA</i>	1,503	1,141	409	55	13.5	31	42	0.1408
<i>rpoA</i>	939	814	438	29	6.6	22	42	ND
<i>pheS</i>	2,533	618	397	46	11.6	30	ND	ND
<i>pepN</i>	1,023	482	411	65	15.8	38	37	0.0741
<i>bcaT</i>	1,047	493	315	87	27.6	39	39	0.0738
<i>pepX</i>	2,269	602	422	81	19.3	26	39	0.0806
SSU rRNA	4,050	650	420	14	3.3	15	ND	ND

^a This subset, apart from *pheS*, does not comprise any ambiguous nucleotides.

^b Pairwise ratios of nonsynonymous to synonymous substitutions (dN/dS) in sequences were calculated by using the method of Nei and Gojobori (25). ND, not determined.

GGGA of the other *L. lactis* subspecies and *Lactococcus* species. A third minor subgroup C is composed of *L. lactis* subsp. *lactis* strains P7266 (NIZO 2206) and P7304 (NIZO 2207); both were able to grow at 37°C. Biovar diacetylactis strains were not differentiated from other *L. lactis* subsp. *lactis* cultures using partial SSU rRNA gene sequence analysis. Note that MG1363, which is commonly referred to as *L. lactis* subsp. *cremoris*, is actually *L. lactis* subsp. *lactis*, with a genotype like that of the *L. lactis* subsp. *cremoris* type strain (19, 38).

MLSA of *L. lactis* strains. Partial DNA sequences of *atpA*, *rpoA*, *pheS*, *bcaT*, *pepN*, and *pepXP* were obtained for 89 *L. lactis* subsp. *lactis* and *L. lactis* subsp. *cremoris* isolates using the primers and temperatures listed in Table 2. No amplification products were obtained with *L. lactis* subsp. *hordniae* and other *Lactococcus* species. Decreasing the annealing temperature by 4°C resulted in additional, presumably nonspecific, amplicons in some of the isolates, while others still did not yield a PCR product and, therefore, were not used for further analysis. As a consequence, the following results are limited to *L. lactis* subsp. *lactis* and *L. lactis* subsp. *cremoris* diversity only.

A limited number of sequences with nucleotide ambiguities were observed for *bcaT* (five sequences with 1 or 2 ambiguities), *pepN* (seven sequences with up to 15 ambiguities), and *pepXP* (seven sequences with 1 or 2 single nucleotide deletions). All positions containing ambiguities were excluded from further analysis. Moreover, no sequences were obtained of *pepN* in strains KF7 (NIZO 2219) and KF 178 (NIZO 2234), of *pepXP* in strains P7266 and P7304, and of *bcaT* in strains P7266, KF 178, KW2 (NIZO 2247), and AM3 (NIZO 2251). Sequences of these isolates were omitted from the composite MLSA cluster analysis presented in Fig. 2 and Table 3. Cluster analysis of the sequences of *atpA*, *rpoA*, *bcaT*, *pepN*, and *pepXP* individually (85 to 89 isolates) and as a composite data set (five loci for MLSA; 83 isolates) (Fig. 2) revealed two distinct clusters corresponding to subgroups A and B in the cluster analysis based on partial SSU rRNA gene sequences (Fig. 1 and 2). Isolates of SSU rRNA gene sequence subgroup C yielded deviating five-locus MLSA sequences, each forming a single strain cluster, designated C₁ and C₂, supporting a more distant relation to the other isolates in this study. In the five-locus MLSA cluster analysis, subgroup A includes all isolates showing a biovar diacetylactis phenotype cluster together at high similarity (>99%) with isolates from dairy substrates (Fig. 2, 5 loci MLSA). This is in contrast with the partial SSU rRNA gene sequence cluster analysis, where the *L. lactis* subsp. *lactis*

dairy isolates were scattered throughout cluster A (Fig. 2, SSU rRNA gene). Only ATCC 19435 clusters at a greater distance from the other isolates from dairy substrates. The single biovar diacetylactis isolate M20 not of dairy origin also clusters at some distance from most of the isolates from dairy substrates.

The genes involved in flavor formation show a larger sequence variation than the housekeeping genes (Table 3). In particular, sequence variation in *rpoA* is limited and close to the SSU rRNA gene.

The observed sequence variation in the partial *pheS* gene sequence (Fig. 2) is substantial, and neighbor-joining cluster analysis resulted in groupings different from those described above. Again, two subgroups were found, one referred to as B+ comprising the members of cluster B (Fig. 1 and 2) with 26 additional *L. lactis* subsp. *lactis* isolates; the other, A-, comprises 43 *L. lactis* subsp. *lactis* isolates including all biovar diacetylactis isolates (Fig. 2, PheS). Limited unexplained ambiguous nucleotides were found in the *pheS* gene sequences in a pattern shared among (most of) the isolates. These nucleotides were not included in the cluster analysis (Fig. 2, PheS).

(GTG)₅-PCR genomic fingerprint analysis. In addition to the genetic approaches described above, the subspecific diversity of *L. lactis* was analyzed with (GTG)₅-PCR genomic fingerprinting. Product moment-UPGMA cluster analysis revealed two distinct clusters (Fig. 3) highly similar to the subgroups A and B observed with SSU rRNA gene sequence analysis and the composite five-locus MLSA clustering (Fig. 2, SSU rRNA gene sequence and five-locus MLSA). From the minor subgroup C, isolate P7304 is found in cluster A. Within the subgroups little diversity was observed. The results confirm the close relationship of *L. lactis* subsp. *lactis* and *L. lactis* subsp. *lactis* biovar diacetylactis since these strains remained indistinguishable using (GTG)₅-PCR genomic fingerprinting analysis. The phenotypic differences between isolates are apparently not reflected in the genotype as sampled in the current collection by this method. The (GTG)₅-PCR genomic fingerprint profiles of the two *L. lactis* subsp. *hordniae* strains show similarity with group B (data not shown) while partial SSU rRNA gene sequences cluster with group A (Fig. 1).

DISCUSSION

Analysis of a collection of strains from various dairy and plant fermentations and a wide range of geographic locations, including Europe, Asia, New Zealand, and diverse environ-




FIG. 3. Product-moment and UPGMA cluster analysis of (GTG)₅-PCR genomic fingerprint profiles of *L. lactis* isolates. The isolates cluster in two subgroups, cluster A comprising 11 *L. lactis* subsp. *cremoris* and 9 *L. lactis* subsp. *lactis* isolates and cluster B including the other 67 *L. lactis* subsp. *lactis* isolates. Isolates obtained from dairy substrates are indicated (■).

mental niches, provides a comprehensive view of the molecular and phenotypic diversity of the species *L. lactis*.

Three molecular typing methods were applied for analysis of *L. lactis* diversity, which allowed high-resolution analysis of the molecular diversity within the species. This newly developed MLSA typing scheme for *L. lactis* revealed unique sequence types for most strains, allowing high-resolution diversity analysis of the current strain collection. Partial SSU rRNA gene sequence analysis, (GTG)₅-PCR genomic fingerprinting analysis, and a novel MLSA scheme all revealed two major, distinct genomic lineages within the species that are dissimilar to groups defined on the basis on extensive phenotypic analysis, including specific assays applied in the validly published classification of the species (15, 34).

Two of the three genomic lineages found are already supported with early work by Garvie and Farrow (9), DNA-DNA homology studies by Jarvis and Jarvis (15), and more recent genotypic analyses including gene sequence analyses (2, 24, 27, 38). One genomic lineage consists almost exclusively of strains of the *L. lactis* subsp. *lactis* including biovar diacetylactis. The second lineage comprised 10 out of the 11 analyzed *L. lactis* subsp. *cremoris* isolates and 10 isolates with an *L. lactis* subsp. *lactis* phenotype. In the current study 9 *L. lactis* subsp. *lactis* isolates with an *L. lactis* subsp. *cremoris* type-strain-like genotype were identified based on SSU rRNA gene sequence, five-locus MLSA, and (GTG)₅ genomic fingerprinting. A reciprocal condition was not reported by Jarvis and Jarvis (15) and Garvie (10), but it was described by Kelly and Ward (16). In our analysis a clear third genomic lineage can be recognized that, to our knowledge, has not been described before. It consists of two isolates with an *L. lactis* subsp. *lactis* phenotype (P7266 and P7304) isolated from litter on pastures, and their partial SSU rRNA gene sequences (Fig. 1 and 2) and five-locus MLSA sequences (Fig. 2) group separately from the other *L. lactis* isolates. Phenotypic differences, however, are less distinct. The SSU rRNA gene sequences of the strains showed high similarity to Ribosomal Database Project II-derived lactococcal sequences of isolates from freshwater prawn (5), an organism that may be found in litter on pastures.

One of the six MLSA gene targets, *pheS*, showed two major groups, different from those revealed by the other genotyping methods and not reported before. One group is formed by the *L. lactis* subsp. *cremoris* type-strain-like genotype (based on SSU rRNA gene sequence and five-locus MLSA) amended with a large portion of *L. lactis* subsp. *lactis* isolates. The other group comprised the remaining *L. lactis* subsp. *lactis* isolates. Congruence of a *pheS* genotype with another genotype or phenotype was not observed. This phenomenon cannot be explained by a single horizontal gene transfer event, as has been identified with other genes from comparative genome analysis of *L. lactis* (2), nor can it be explained by gene duplication. Gene duplication of *pheS* was not found in strains IL-1403, SK11, and MG1363, for which full genome sequences have been published (3, 22, 40).

L. lactis subsp. *lactis* and *L. lactis* subsp. *cremoris* are defined based on their phenotypes (34). In general, the isolates with an *L. lactis* subsp. *lactis* phenotype showed a partial SSU rRNA gene sequence similar to the *L. lactis* subsp. *lactis* type strain while a limited number of isolates showed an *L. lactis* subsp. *cremoris*-(type strain)-like genotype. Unfortunately, this has

been turned around by several groups using a subspecific classification based on genotype, causing confusion in the taxonomy of *L. lactis*. This started with the application of supposedly subspecies-specific probes (18, 31, 32) and continued with additional methods for genetic differentiation such as SSU rRNA gene sequencing (39) and other genotyping methods (see reference 38 and references therein). However, based on our findings, there seems to be no basis for redefinition of the *L. lactis* subspecies. The phenotypic differences listed in the current classification are sufficient to distinguish the subspecies (34). However, one should be aware that the subspecific identification of *L. lactis* in the literature from the 1990s to the present is often not based on the phenotype and may therefore be confusing. This confusion can be avoided by using the current phenotypic classification amended with a "type strain-like-genotype" classification, which will provide a direct subspecific phylogenetic reference.

The remarkable taxonomic structure of *L. lactis* is also evident from the genome sequences of three different strains that have been published (3, 40). These strains include well-known model strains of dairy origin such as strain IL-1403 (*L. lactis* subsp. *lactis* type-strain-like genotype and *L. lactis* subsp. *lactis* phenotype), strain MG363 (*L. lactis* subsp. *cremoris* type-strain-like genotype and *L. lactis* subsp. *lactis* phenotype), and strain SK11 (*L. lactis* subsp. *cremoris* type-strain-like genotype and *L. lactis* subsp. *cremoris* phenotype). Strain IL-1403 is actually a plasmid-cured derivative of the biovar diacetylactis strain CNRZ157 rendering it *cit* negative. This analysis and the reports of other investigators (27, 38) show that these three strains clearly represent only a small part of the diversity present within this subspecies. It can be expected that the adaptation to diverse plant substrates of the nondairy strains has resulted in the development of unique traits and phenotypes that can be utilized in dairy and other food fermenting applications. Recently, we have sequenced the genomes of two plant isolates (38). Further analysis of these data and genotyping with DNA microarrays in our laboratory will provide insight into the lack of congruence of the *pheS* genotype with other features and will allow a deeper understanding of the genomic and functional diversity within the species.

ACKNOWLEDGMENTS

We acknowledge Jan Hoolwerf and Arjen Wagendorp for their technical assistance. The following people are acknowledged for kindly providing strains used in this study: Andreas Ulrich and Thomas Müller, Centre for Agricultural Landscape and Land Use Research Münchenberg Institute of Microbial Ecology and Soil Biology Münchenberg, Germany; Sang-Hee Park, Laboratory of Veterinary Public Health Graduate School of Agricultural and Life Sciences, the University of Tokyo, Japan; and U. Shillinger, Federal Research Centre for Nutrition Institute of Hygiene and Toxicology Karlsruhe, Germany. Meike te Giffel and Giovanna Felis are acknowledged for valuable comments on the manuscript and Bart Hoste and Marc Vancanneyt for useful discussions including *Lactococcus* taxonomy.

D.G. and J.S. are indebted to the Fund for Scientific Research, Flanders (Belgium), for a position as a postdoctoral fellow and research grants, respectively.

REFERENCES

1. Ayad, E. H., A. Verheul, W. J. Engels, J. T. Wouters, and G. Smit. 2001. Enhanced flavour formation by combination of selected lactococci from industrial and artisanal origin with focus on completion of a metabolic pathway. *J. Appl. Microbiol.* 90:59–67.

2. Bolotin, A., B. Quinquis, A. Sorokin, and D. S. Ehrlich. 2004. Recent genetic transfer between *Lactococcus lactis* and enterobacteria. *J. Bacteriol.* **186**: 6671–6677.
3. Bolotin, A., P. Wincker, S. Mauger, O. Jaillon, K. Malarme, J. Weissenbach, S. Ehrlich, and A. Sorokin. 2001. The complete genome sequence of the lactic acid bacterium *Lactococcus lactis* ssp. *lactis* IL1403. *Genome Res.* **11**:731–753.
4. Bougnoux, M. E., D. Diogo, N. Francois, B. Sendid, S. Veirmeire, J. F. Colombe, C. Bouchier, H. Van Kruiningen, C. d'Enfert, and D. Poulaire. 2006. Multilocus sequence typing reveals intrafamilial transmission and microevolutions of *Candida albicans* isolates from the human digestive tract. *J. Clin. Microbiol.* **44**:1810–1820.
5. Cole, J. R., B. Chai, R. J. Farris, Q. Wang, S. A. Kulam, D. M. McGarrell, G. M. Garrity, and J. M. Tiedje. 2005. The Ribosomal Database Project (RDP-II): sequences and tools for high-throughput rRNA analysis. *Nucleic Acids Res.* **33**:D294–D296.
6. de Las Rivas, B., A. Marcobal, and R. Munoz. 2006. Development of a multilocus sequence typing method for analysis of *Lactobacillus plantarum* strains. *Microbiology* **152**:85–93.
7. Erlandson, K., and C. A. Batt. 1997. Strain-specific differentiation of lactococci in mixed starter culture populations using randomly amplified polymorphic DNA-derived probes. *Appl. Environ. Microbiol.* **63**:2702–2707.
8. Galesloot, T. E., I. Hassing, and J. Stadhouders. 1961. Agar media for the isolation and enumeration of aroma bacteria in starters. *Neth. Milk Dairy J.* **15**:145–150.
9. Garvie, E., and J. Farrow. 1982. *Streptococcus lactis* subsp. *cremoris* (Orla-Jensen) comb. nov. and *Streptococcus lactis* subsp. *diacetilactis* (Matuszewski et al.) nom. rev., comb. nov. *Int. J. Syst. Bacteriol.* **32**:453–455.
10. Garvie, E. I. 1978. *Streptococcus raffinolactis* (Orla-Jensen and Hansen): a group N streptococcus found in raw milk. *Int. J. Syst. Bacteriol.* **28**:190–193.
11. Gevers, D., G. Huys, and J. Swings. 2001. Applicability of rep-PCR fingerprinting for identification of *Lactobacillus* species. *FEMS Microbiol. Lett.* **205**:31–36.
12. Helgason, E., N. J. Tourasse, R. Meisal, D. A. Caugant, and A. B. Kolsto. 2004. Multilocus sequence typing scheme for bacteria of the *Bacillus cereus* group. *Appl. Environ. Microbiol.* **70**:191–201.
13. Holt, J. G., N. R. Krieg, P. H. A. Sneath, J. T. Staley, and S. T. Williams. 1994. Bergey's manual of determinative bacteriology, 9th ed. Williams and Wilkins Co., Baltimore, MD.
14. Hommais, F., S. Pereira, C. Acquaviva, P. Escobar-Paramo, and E. Denamur. 2005. Single-nucleotide polymorphism phytotyping of *Escherichia coli*. *Appl. Environ. Microbiol.* **71**:4784–4792.
15. Jarvis, A. W., and B. D. Jarvis. 1981. Deoxyribonucleic acid homology among lactic streptococci. *Appl. Environ. Microbiol.* **41**:77–83.
16. Kelly, W., and L. Ward. 2002. Genotypic vs. phenotypic biodiversity in *Lactococcus lactis*. *Microbiology* **148**:3332–3333.
17. Kelly, W. J., G. P. Davey, and L. J. Ward. 1998. Characterization of lactococci isolated from minimally processed fresh fruit and vegetables. *Int. J. Food Microbiol.* **45**:85–92.
18. Klijn, N., A. H. Weerkamp, and W. M. de Vos. 1995. Detection and characterization of lactose-utilizing *Lactococcus* spp. in natural ecosystems. *Appl. Environ. Microbiol.* **61**:788–792.
19. Kok, J., G. Buist, A. L. Zomer, S. A. van Huijum, and O. P. Kuipers. 2005. Comparative and functional genomics of lactococci. *FEMS Microbiol. Rev.* **29**:411–433.
20. Liu, M., F. H. van Enckevort, and R. J. Siezen. 2005. Genome update: lactic acid bacteria genome sequencing is booming. *Microbiology* **151**:3811–3814.
21. Maiden, M. C., J. A. Bygraves, E. Feil, G. Morelli, J. E. Russell, R. Urwin, Q. Zhang, J. Zhou, K. Zurth, D. A. Caugant, I. M. Feavers, M. Achtman, and B. G. Spratt. 1998. Multilocus sequence typing: a portable approach to the identification of clones within populations of pathogenic microorganisms. *Proc. Natl. Acad. Sci. USA* **95**:3140–3145.
22. Makarova, K., A. Slesarev, Y. Wolf, A. Sorokin, B. Mirkin, E. Koonin, A. Pavlov, N. Pavlova, V. Karamychev, N. Polouchine, V. Shakhova, I. Grigoriev, Y. Lou, D. Rohksar, S. Lucas, K. Huang, D. M. Goodstein, T. Hawkinse, V. Plengvidhya, D. Welker, J. Hughes, Y. Goh, A. Benson, K. Baldwin, J.-H. Lee, I. Diaz-Muniz, B. Dosti, V. Smeianov, W. Wechter, R. Barabote, G. Lorca, E. Altermann, R. Barrangou, B. Ganesan, Y. Xie, H. Rawsthorne, D. Tamir, C. Parker, F. Breidt, J. Broadbent, R. Hutchins, D. O'Sullivan, J. Steele, G. Unlu, M. Saier, T. Klaenhammer, P. Richardson, S. Kozyavkin, B. Weimerd, and D. Mills. 2006. Comparative genomics of the lactic acid bacteria. *Proc. Natl. Acad. Sci. USA* **103**:15611–15616.
23. Mangin, I., D. Corroler, A. Reinhardt, and M. Gueguen. 1999. Genetic diversity among dairy lactococcal strains investigated by polymerase chain reaction with three arbitrary primers. *J. Appl. Microbiol.* **86**:514–520.
24. Mori, S., K. Mori, I. Suzuki, and T. Kasumi. 2004. Phylogenetic analysis of *Lactococcus lactis* subspecies based on decoding the sequence of the *pepT* tripeptidase gene, the *pepV* dipeptidase gene and 16S rRNA. *Syst. Appl. Microbiol.* **27**:414–422.
25. Nei, M., and T. Gojobori. 1986. Simple methods for estimating the numbers of synonymous and nonsynonymous nucleotide substitutions. *Mol. Biol. Evol.* **3**:418–426.
26. Niven, C. F., K. L. Smiley, and J. M. Sherman. 1942. The hydrolysis of arginine by streptococci. *J. Bacteriol.* **43**:651–660.
27. Nomura, M., M. Kobayashi, T. Narita, H. Kimoto-Nira, and T. Okamoto. 2006. Phenotypic and molecular characterization of *Lactococcus lactis* from milk and plants. *J. Appl. Microbiol.* **101**:396–405.
28. Pu, Z. Y., M. Dobos, G. K. Limsowtin, and I. B. Powell. 2002. Integrated polymerase chain reaction-based procedures for the detection and identification of species and subspecies of the gram-positive bacterial genus *Lactococcus*. *J. Appl. Microbiol.* **93**:353–361.
29. Rademaker, J. L. W., and F. J. deBruijn. 2004. Computer-assisted pattern analysis of electrophoretic fingerprints and database construction. Chapter 1.7.5, p. 1–50. In G. Kowalchuk, F. deBruijn, I. Head, A. Akkermans, and J. van Elsas (ed.), *Molecular microbial ecology manual*, 2nd ed. Kluwer Academic Publishers, Dordrecht, The Netherlands.
30. Rademaker, J. L. W., F. J. Louws, J. Versalovic, and F. J. deBruijn. 2004. Characterization of the diversity of ecological important microbes by rep-PCR genomic fingerprinting. Chapter 5.3.2, p. 1–33. In G. Kowalchuk, F. deBruijn, I. Head, A. Akkermans, and J. van Elsas (ed.), *Molecular microbial ecology manual*, 2nd ed. Kluwer Academic Publishers, Dordrecht, The Netherlands.
31. Salama, M., W. Sandine, and S. Giovannoni. 1991. Development and application of oligonucleotide probes for identification of *Lactococcus lactis* subsp. *cremoris*. *Appl. Environ. Microbiol.* **57**:1313–1318.
32. Salama, M. S., W. E. Sandine, and S. J. Giovannoni. 1993. Isolation of *Lactococcus lactis* subsp. *cremoris* from nature by colony hybridization with rRNA probes. *Appl. Environ. Microbiol.* **59**:3941–3945.
33. Salcedo, C., L. Arreaza, B. Alcala, L. de la Fuente, and J. A. Vazquez. 2003. Development of a multilocus sequence typing method for analysis of *Listeria monocytogenes* clones. *J. Clin. Microbiol.* **41**:757–762.
34. Schleifer, K. H., J. Kraus, C. Dvorak, R. Kilpper-Balz, M. D. Collins, and W. Fisher. 1985. Transfer of *Streptococcus lactis* and related streptococci to the genus *Lactococcus* gen. nov. *Syst. Appl. Microbiol.* **6**:183–195.
35. Smit, G., B. A. Smit, and W. J. Engels. 2005. Flavour formation by lactic acid bacteria and biochemical flavour profiling of cheese products. *FEMS Microbiol. Rev.* **29**:591–610.
36. Tailliez, P., J. Tremblay, S. D. Ehrlich, and A. Chopin. 1998. Molecular diversity and relationship within *Lactococcus lactis*, as revealed by randomly amplified polymorphic DNA (RAPD). *Syst. Appl. Microbiol.* **21**:530–538.
37. Torpahl, M., M. N. Skov, D. Sandvang, and D. L. Baggerup. 2005. Genotypic characterization of *Salmonella* by multilocus sequence typing, pulsed-field gel electrophoresis and amplified fragment length polymorphism. *J. Microbiol. Methods* **63**:173–184.
38. van Hylckama Vlieg, J. E., J. L. Rademaker, H. Bachmann, D. Molenaar, W. J. Kelly, and R. J. Siezen. 2006. Natural diversity and adaptive responses of *Lactococcus lactis*. *Curr. Opin. Biotechnol.* **17**:183–190.
39. Ward, L. J., J. C. Brown, and G. P. Davey. 1998. Two methods for the genetic differentiation of *Lactococcus lactis* ssp. *lactis* and *cremoris* based on differences in the 16S rRNA gene sequence. *FEMS Microbiol. Lett.* **166**:15–20.
40. Wegmann, U., M. O'Connell-Motherway, A. Zomer, G. Buist, C. Shearman, C. Canchaya, M. Ventura, A. Goessmann, M. Gasson, O. Kuipers, D. v. Sinderen, and J. Kok. 2007. Complete genome sequence of the prototype lactic acid bacterium *Lactococcus lactis* subsp. *cremoris* MG1363. *J. Bacteriol.* **189**:3256–3270.