

Dichlorido(2,9-dimethyl-1,10-phenanthroline- κ^2N,N')mercury(II)

Ismail Warad,^a Mousa Al-Noaimi,^{b*} Salim F. Haddad^c and Rema Othman^d

^aDepartment of Chemistry, AN-Najah National University, Nablus, Palestinian Territories, ^bDepartment of Chemistry, Hashemite University, Zarqa 13115, Jordan, ^cDepartment of Chemistry, The University of Jordan, Amman 11942, Jordan, and ^dLanguage Centre, Hashemite University, Zarqa 13115, Jordan
Correspondence e-mail: manoaimi@hu.edu.jo


Received 2 January 2013; accepted 11 January 2013

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.016$ Å; R factor = 0.051; wR factor = 0.128; data-to-parameter ratio = 14.7.

The title compound, $[\text{HgCl}_2(\text{C}_{14}\text{H}_{12}\text{N}_2)]$, consists of one 2,9-dimethyl-1,10-phenanthroline (dmphen) ligand chelating the Hg^{II} ion and two chloride ligands coordinating to the Hg^{II} ion, forming a distorted tetrahedral environment. The dmphen ligand is nearly planar (r.m.s. deviation = 0.0225 Å). The dihedral angle between the normal to the plane defined by the Hg^{II} atom and the two Cl atoms and the normal to the plane of the dmphen ring is 81.8 (1)°.

Related literature

For related structures, see Alizadeh (2009); Alizadeh *et al.* (2009); Wang & Zhong (2009); Warad *et al.* (2011). For properties and application of mercury(II) complexes, see: Ramazani *et al.* (2005); Mahjoub *et al.* (2004); Canty & Maker (1976); Canty & Lee (1982).



Experimental

Crystal data

| | |
|---|---|
| $[\text{HgCl}_2(\text{C}_{14}\text{H}_{12}\text{N}_2)]$ | $V = 1462.8(4) \text{ \AA}^3$ |
| $M_r = 479.75$ | $Z = 4$ |
| Monoclinic, $P2_1/c$ | Mo $K\alpha$ radiation |
| $a = 7.5732(13) \text{ \AA}$ | $\mu = 10.87 \text{ mm}^{-1}$ |
| $b = 10.3733(16) \text{ \AA}$ | $T = 293 \text{ K}$ |
| $c = 18.673(2) \text{ \AA}$ | $0.22 \times 0.20 \times 0.18 \text{ mm}$ |
| $\beta = 94.308(12)^\circ$ | |

Data collection

| | |
|---|--|
| Agilent Xcalibur Eos diffractometer | 5483 measured reflections |
| Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Agilent, 2011) | 2564 independent reflections |
| $T_{\text{min}} = 0.106$, $T_{\text{max}} = 0.140$ | 1758 reflections with $I > 2\sigma(I)$ |
| | $R_{\text{int}} = 0.061$ |

Refinement

| | |
|---------------------------------|--|
| $R[F^2 > 2\sigma(F^2)] = 0.051$ | 174 parameters |
| $wR(F^2) = 0.128$ | H-atom parameters constrained |
| $S = 0.99$ | $\Delta\rho_{\text{max}} = 1.81 \text{ e \AA}^{-3}$ |
| 2564 reflections | $\Delta\rho_{\text{min}} = -1.83 \text{ e \AA}^{-3}$ |

Data collection: *CrysAlis PRO* (Agilent, 2011); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP III* (Burnett & Johnson, 1996); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

The project was supported by King Saud University (KSA) and Hashemite University (Jordan). The X-ray structural work was done at Hamdi Mango Center for Scientific Research at The University of Jordan, Amman 11942, Jordan.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BR2220).

References

- Agilent (2011). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
 Alizadeh, R. (2009). *Acta Cryst.* **E65**, m817–m818.
 Alizadeh, R., Heidari, A., Ahmadi, R. & Amani, V. (2009). *Acta Cryst.* **E65**, m483–m484.
 Burnett, M. N. & Johnson, C. K. (1996). *ORTEP III*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
 Canty, A. J. & Lee, C. V. (1982). *Organometallics*, **1**, 1063–1066.
 Canty, A. J. & Maker, A. (1976). *Inorg. Chem.* **15**, 425–430.
 Mahjoub, A., Morsali, A. & Nejad, R. (2004). *Z. Naturforsch. Teil B*, **59**, 1109–1113.
 Ramazani, A., Morsali, A., Dolatyari, L. & Ganjeie, B. (2005). *Z. Naturforsch. Teil B*, **60**, 289–293.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Wang, B. S. & Zhong, H. (2009). *Acta Cryst.* **E65**, m1156.
 Warad, I., Boshala, A., Al-Resayes, S. I., Al-Deyab, S. S. & Rzaigui, M. (2011). *Acta Cryst.* **E67**, m1650.

supplementary materials

Acta Cryst. (2013). E69, m109 [doi:10.1107/S1600536813001086]

Dichlorido(2,9-dimethyl-1,10-phenanthroline- κ^2N,N')mercury(II)

Ismail Warad, Mousa Al-Noaimi, Salim F. Haddad and Rema Othman

Comment

The coordination chemistry of mercury(II) with N-donor ligands is of interest due to applications as solid-state materials (Ramazani *et al.*, 2005; Mahjoub *et al.*, 2004). Hg(II) complexes with bidentate ligands have been obtained in which Hg(II) adopts higher coordination numbers such as complexes of 1,10-phenanthroline (Canty & Maker, 1976) and N-substituted pyrazole (Canty & Lee, 1982). The molecular structure of [HgCl₂(C₁₄H₁₂N₂)], along with the numbering scheme is shown in Fig. 1. HgCl₂ is chelated by the bidentate phenanthroline molecule and that the coordination of the nitrogen and chlorine atoms about the Hg atom is essentially a distorted tetrahedral environment (Fig. 1).

Experimental


The desired complex was prepared by mixing of mercury chloride (HgCl₂, 39.7 mg, 0.14 mmol) in methanol (10 ml) with dmphen (32.0 mg, 0.15 mmol) in dichloromethane (5 ml) is stirred for one hour at room temperature. The obtained solution was concentrated to about 2 ml under reduced pressure and mixed to 30 ml of diethyl ether. The white precipitate was filtered and dried. Suitable colourless crystals were obtained by slow diffusion of diethyl ether into a solution of the complex in dichloromethane.

Refinement

All nonhydrogen atoms were refined anisotropically. H atoms were positioned geometrically, with C-H = 0.93 and 0.96 Å for aromatic and methyl H, respectively, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. Highest difference peak and hole are 1.81 and -1.83 e/Å³ close to the Hg atom.

Computing details

Data collection: *CrysAlis PRO* (Agilent, 2011); cell refinement: *CrysAlis PRO* (Agilent, 2011); data reduction: *CrysAlis PRO* (Agilent, 2011); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP III* (Burnett & Johnson, 1996); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).


Figure 1

An ORTEP (Burnett & Johnson, 1996) view of $\text{Hg}(\text{Cl})_2(\text{dmphen})$. Displacement ellipsoids are drawn at the 30% probability level. H atoms are represented as small spheres of arbitrary radii.

Dichlorido(2,9-dimethyl-1,10-phenanthroline- κ^2N,N')mercury(II)

Crystal data

$[\text{HgCl}_2(\text{C}_{14}\text{H}_{12}\text{N}_2)]$

$M_r = 479.75$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2_1/c$

$a = 7.5732$ (13) Å

$b = 10.3733$ (16) Å

$c = 18.673$ (2) Å

$\beta = 94.308$ (12)°

$V = 1462.8$ (4) Å³

$Z = 4$

$F(000) = 896$

$D_x = 2.178$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1534 reflections

$\theta = 2.9\text{--}29.0^\circ$

$\mu = 10.87$ mm⁻¹

$T = 293$ K

Block, colourless

$0.22 \times 0.20 \times 0.18$ mm

Data collection

Agilent Xcalibur Eos

diffractometer

Radiation source: Enhance (Mo) X-ray Source

Graphite monochromator

Detector resolution: 16.0534 pixels mm⁻¹

ω scans

Absorption correction: multi-scan

(*CrysAlis PRO*; Agilent, 2011)

$T_{\min} = 0.106$, $T_{\max} = 0.140$

5483 measured reflections

2564 independent reflections
 1758 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.061$
 $\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 2.9^\circ$

$h = -9 \rightarrow 7$
 $k = -12 \rightarrow 12$
 $l = -16 \rightarrow 22$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.051$
 $wR(F^2) = 0.128$
 $S = 0.99$
 2564 reflections
 174 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0521P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 1.81 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -1.83 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | x | y | z | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|------|-------------|--------------|--------------|----------------------------------|
| Hg1 | 0.21008 (6) | 0.28289 (4) | 0.39470 (2) | 0.0597 (2) |
| Cl1 | -0.0404 (4) | 0.3290 (4) | 0.31096 (16) | 0.0818 (10) |
| Cl2 | 0.4266 (5) | 0.4523 (3) | 0.39503 (19) | 0.0871 (11) |
| C5 | 0.3385 (16) | -0.2144 (11) | 0.4986 (7) | 0.063 (3) |
| H5A | 0.3736 | -0.3002 | 0.4975 | 0.076* |
| N1 | 0.2952 (11) | 0.0704 (7) | 0.3787 (4) | 0.047 (2) |
| C1 | 0.3465 (14) | 0.0196 (11) | 0.3185 (6) | 0.057 (3) |
| C10 | 0.1290 (15) | 0.2260 (10) | 0.5625 (7) | 0.055 (3) |
| C11 | 0.2939 (12) | -0.0049 (11) | 0.4375 (5) | 0.048 (2) |
| C3 | 0.3905 (15) | -0.1866 (12) | 0.3704 (7) | 0.060 (3) |
| H3A | 0.4210 | -0.2731 | 0.3670 | 0.072* |
| C8 | 0.1676 (15) | 0.0244 (12) | 0.6240 (6) | 0.067 (3) |
| H8A | 0.1636 | -0.0247 | 0.6655 | 0.080* |
| C12 | 0.2350 (11) | 0.0470 (11) | 0.5026 (5) | 0.048 (3) |
| N2 | 0.1849 (11) | 0.1747 (8) | 0.5019 (4) | 0.045 (2) |
| C7 | 0.2276 (13) | -0.0328 (10) | 0.5619 (6) | 0.052 (3) |
| C14 | 0.0775 (16) | 0.3651 (12) | 0.5595 (6) | 0.073 (4) |
| H14A | 0.1822 | 0.4176 | 0.5615 | 0.110* |
| H14B | 0.0100 | 0.3853 | 0.5995 | 0.110* |
| H14C | 0.0073 | 0.3819 | 0.5155 | 0.110* |
| C4 | 0.3430 (13) | -0.1380 (10) | 0.4363 (6) | 0.052 (3) |
| C2 | 0.3929 (15) | -0.1104 (11) | 0.3118 (7) | 0.066 (3) |

| | | | | |
|------|-------------|--------------|------------|-----------|
| H2A | 0.4245 | -0.1434 | 0.2683 | 0.080* |
| C6 | 0.2834 (14) | -0.1637 (11) | 0.5598 (7) | 0.062 (3) |
| H6A | 0.2820 | -0.2147 | 0.6007 | 0.074* |
| C9 | 0.1155 (14) | 0.1497 (14) | 0.6246 (5) | 0.061 (3) |
| H9A | 0.0715 | 0.1850 | 0.6654 | 0.074* |
| C13 | 0.3475 (18) | 0.1076 (12) | 0.2547 (6) | 0.083 (4) |
| H13A | 0.3449 | 0.0572 | 0.2116 | 0.124* |
| H13B | 0.4528 | 0.1594 | 0.2587 | 0.124* |
| H13C | 0.2453 | 0.1626 | 0.2532 | 0.124* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|------------|-------------|-------------|--------------|-------------|--------------|
| Hg1 | 0.0813 (4) | 0.0419 (4) | 0.0565 (3) | 0.0057 (2) | 0.0083 (3) | 0.00836 (19) |
| Cl1 | 0.075 (2) | 0.111 (3) | 0.0590 (19) | 0.016 (2) | 0.0022 (15) | 0.0250 (18) |
| Cl2 | 0.106 (3) | 0.0474 (18) | 0.107 (3) | -0.0145 (19) | 0.003 (2) | 0.0152 (18) |
| C5 | 0.070 (8) | 0.046 (8) | 0.075 (9) | -0.002 (6) | 0.009 (6) | 0.023 (7) |
| N1 | 0.056 (5) | 0.033 (5) | 0.051 (5) | -0.001 (4) | 0.002 (4) | -0.003 (4) |
| C1 | 0.061 (7) | 0.052 (7) | 0.059 (7) | 0.003 (6) | 0.014 (5) | 0.003 (6) |
| C10 | 0.053 (7) | 0.047 (7) | 0.065 (8) | 0.003 (5) | 0.007 (5) | -0.008 (6) |
| C11 | 0.037 (6) | 0.056 (7) | 0.049 (6) | -0.003 (5) | -0.001 (4) | 0.003 (6) |
| C3 | 0.050 (7) | 0.049 (7) | 0.080 (9) | 0.007 (5) | -0.004 (6) | -0.003 (7) |
| C8 | 0.071 (8) | 0.066 (8) | 0.062 (8) | -0.016 (7) | -0.002 (6) | 0.017 (7) |
| C12 | 0.020 (5) | 0.060 (7) | 0.062 (7) | -0.006 (5) | -0.007 (4) | 0.008 (6) |
| N2 | 0.051 (5) | 0.043 (5) | 0.041 (5) | -0.006 (4) | 0.004 (4) | -0.002 (4) |
| C7 | 0.062 (7) | 0.038 (6) | 0.055 (7) | -0.007 (5) | 0.001 (5) | 0.015 (5) |
| C14 | 0.078 (9) | 0.088 (10) | 0.055 (7) | 0.002 (8) | 0.015 (6) | -0.021 (7) |
| C4 | 0.042 (6) | 0.036 (6) | 0.077 (8) | -0.007 (5) | 0.000 (5) | 0.005 (6) |
| C2 | 0.076 (9) | 0.057 (8) | 0.065 (8) | 0.015 (7) | -0.001 (6) | -0.016 (7) |
| C6 | 0.048 (7) | 0.060 (8) | 0.075 (9) | -0.014 (6) | -0.006 (6) | 0.030 (7) |
| C9 | 0.051 (7) | 0.099 (10) | 0.035 (6) | -0.009 (7) | 0.007 (4) | 0.005 (7) |
| C13 | 0.133 (13) | 0.059 (8) | 0.061 (8) | 0.001 (8) | 0.031 (8) | 0.006 (7) |

Geometric parameters (\AA , $^\circ$)

| | | | |
|---------|------------|----------|------------|
| Hg1—N2 | 2.314 (8) | C3—C4 | 1.401 (15) |
| Hg1—N1 | 2.322 (8) | C3—H3A | 0.9300 |
| Hg1—Cl2 | 2.403 (3) | C8—C9 | 1.359 (16) |
| Hg1—Cl1 | 2.414 (3) | C8—C7 | 1.408 (15) |
| C5—C6 | 1.352 (16) | C8—H8A | 0.9300 |
| C5—C4 | 1.410 (15) | C12—N2 | 1.378 (13) |
| C5—H5A | 0.9300 | C12—C7 | 1.386 (13) |
| N1—C1 | 1.326 (12) | C7—C6 | 1.424 (15) |
| N1—C11 | 1.348 (12) | C14—H14A | 0.9600 |
| C1—C2 | 1.402 (15) | C14—H14B | 0.9600 |
| C1—C13 | 1.501 (15) | C14—H14C | 0.9600 |
| C10—N2 | 1.348 (13) | C2—H2A | 0.9300 |
| C10—C9 | 1.414 (15) | C6—H6A | 0.9300 |
| C10—C14 | 1.494 (15) | C9—H9A | 0.9300 |
| C11—C4 | 1.430 (14) | C13—H13A | 0.9600 |

| | | | |
|-------------|-------------|---------------|------------|
| C11—C12 | 1.432 (13) | C13—H13B | 0.9600 |
| C3—C2 | 1.350 (16) | C13—H13C | 0.9600 |
| N2—Hg1—N1 | 72.1 (3) | C10—N2—C12 | 118.3 (9) |
| N2—Hg1—C12 | 116.9 (2) | C10—N2—Hg1 | 125.9 (7) |
| N1—Hg1—C12 | 119.9 (2) | C12—N2—Hg1 | 115.8 (6) |
| N2—Hg1—C11 | 123.0 (2) | C12—C7—C8 | 116.1 (10) |
| N1—Hg1—C11 | 108.5 (2) | C12—C7—C6 | 121.2 (11) |
| C12—Hg1—C11 | 111.07 (13) | C8—C7—C6 | 122.6 (10) |
| C6—C5—C4 | 120.5 (11) | C10—C14—H14A | 109.5 |
| C6—C5—H5A | 119.8 | C10—C14—H14B | 109.5 |
| C4—C5—H5A | 119.8 | H14A—C14—H14B | 109.5 |
| C1—N1—C11 | 118.8 (9) | C10—C14—H14C | 109.5 |
| C1—N1—Hg1 | 126.0 (7) | H14A—C14—H14C | 109.5 |
| C11—N1—Hg1 | 115.2 (7) | H14B—C14—H14C | 109.5 |
| N1—C1—C2 | 123.3 (10) | C3—C4—C5 | 123.1 (10) |
| N1—C1—C13 | 116.8 (10) | C3—C4—C11 | 116.5 (10) |
| C2—C1—C13 | 119.9 (10) | C5—C4—C11 | 120.4 (10) |
| N2—C10—C9 | 120.9 (10) | C3—C2—C1 | 118.2 (11) |
| N2—C10—C14 | 116.5 (10) | C3—C2—H2A | 120.9 |
| C9—C10—C14 | 122.5 (10) | C1—C2—H2A | 120.9 |
| N1—C11—C4 | 121.9 (9) | C5—C6—C7 | 120.3 (11) |
| N1—C11—C12 | 119.7 (10) | C5—C6—H6A | 119.8 |
| C4—C11—C12 | 118.4 (10) | C7—C6—H6A | 119.8 |
| C2—C3—C4 | 121.4 (11) | C8—C9—C10 | 119.3 (11) |
| C2—C3—H3A | 119.3 | C8—C9—H9A | 120.3 |
| C4—C3—H3A | 119.3 | C10—C9—H9A | 120.3 |
| C9—C8—C7 | 121.5 (11) | C1—C13—H13A | 109.5 |
| C9—C8—H8A | 119.3 | C1—C13—H13B | 109.5 |
| C7—C8—H8A | 119.3 | H13A—C13—H13B | 109.5 |
| N2—C12—C7 | 123.7 (10) | C1—C13—H13C | 109.5 |
| N2—C12—C11 | 117.1 (9) | H13A—C13—H13C | 109.5 |
| C7—C12—C11 | 119.1 (10) | H13B—C13—H13C | 109.5 |